

Impacts of Cheatgrass on Mammal, Bird, and Butterfly Populations in a Rocky Mountain Foothills Grassland

Anyll Markevich, Stephen R. Jones, Christel Markevich

Report for Boulder County Parks and Open Space

12/06/2020

Executive Summary

Impacts of Cheatgrass on Mammal, Bird, and Butterfly Populations in a Rocky Mountain Foothills

Grassland. Anyll Markevich (Primary Investigator: anyllmarkevich@gmail.com, 303-442-4475), Stephen

R. Jones (Grant Recipient: curlwsj@comcast.net, 303-494-2468), Christel Markevich. 12/07/2020.

Understanding the influence of cheatgrass on mammal, bird, and butterfly populations is vital for proper wildlife management, especially as cheatgrass continues to spread across the western United States. Few studies have investigated these relationships. This study aims to give a preliminary indication of the effects of cheatgrass on wildlife populations of wildlife populations within a Rocky Mountain foothills grassland..

We located eight grassland plots, varying in cheatgrass cover, at the base of the Rocky Mountain Front Range south of Lyons, Colorado. Remote triggering cameras, point counts, and transect counts were used to quantify numbers of mammals, birds, and butterflies respectively. We used a combination of ANOVA analysis and Linear Correlation Analysis to analyze our data.

We found that mammals were less numerous in areas with high cheatgrass cover. We did not find a statistically significant relationship between birds and cheatgrass, through birds numbers tended to decrease with increased cheatgrass cover. Butterflies and butterfly species richness (total number of species) showed statistically significant negative correlation with cheatgrass.

Our analysis suggested that mammals and birds are unaffected by cheatgrass up to a certain threshold of cheatgrass cover above which they are repelled. Conversely, butterflies were linearly correlated with cheatgrass cover. The ability of mammals and birds to move across relatively large distances may allow

27 them to find enough food even in areas interspersed with a certain amount of cheatgrass. Meanwhile the
28 small home ranges of butterflies and their extreme dependance on local nectar sources may cause
29 butterfly numbers to be reduced in tandem with the reduced availability of nectar sources in places where
30 cheatgrass outcompetes the local vegetation.

31
32 Although our study does not prove causal relationships, we believe that the differences in animal
33 populations among plots were in fact driven by cheatgrass infestation. Our belief is supported by the
34 generally weak relationships we observed between animal populations and plot characteristics other than
35 cheatgrass cover.

36
37 Future research should investigate the impact on wildlife of different treatment methods to control
38 cheatgrass. We recommend that such research compare wildlife populations over appropriately long
39 timescales on untreated plots relatively free of cheatgrass, on untreated plots infested by cheatgrass, and
40 on previously infested treated plots. The standard for treatment success should include restoration to
41 conditions similar to those found in untreated, relatively cheatgrass free areas, not just improvement over
42 untreated cheatgrass infested areas.

43
44 Additional possible management implications of this study include:

- 45 • Agencies funding research investigating which cheatgrass removal methods are most beneficial to
46 wildlife (including alternative methods such as soil regeneration)
- 47 • Biologists accounting for cheatgrass infestation when evaluating the quality of wildlife habitats
- 48 • Agencies prioritizing native plant restoration in ecologically important areas, such as breeding
49 grounds

50

51

Abstract

We studied impacts of cheatgrass (*Anisantha* spp.) infestation on mammals, birds, and butterflies in a foothills grassland south of Lyons, Colorado. The impacts of cheatgrass on wildlife have been poorly studied despite the prevalence of cheatgrass infestation in the United States. We established eight, 75 m radius circular plots in areas of mixed-grass prairie (also containing native shrubs and ponderosa pines) on two Boulder County Parks and Open Space properties. Four of the plots were located in areas with relatively low (3-13%) cheatgrass cover, four plots in areas with relatively high (24-35 %) cheatgrass cover. We sampled vegetation within each plot using a Point-intercept with Grid Quadrant method. We installed motion-activated wildlife cameras at the center of each plot. The cameras were active over a 3 month period. We conducted four early-morning point-count bird surveys in each plot between May 29th and July 9th, counting all birds seen or heard perching or foraging within the plot over 8-minute intervals. We counted butterflies within 30 m of 150 m transects bisecting the plots on 6 days between June 6th and August 10th. We found a significant negative relationship between total mammals and cheatgrass cover of plots ($p=.02$) and an insignificant negative relationship between mammal species richness and cheatgrass cover of plots ($p=.15$). While bird abundance and bird species richness was greater on low cheatgrass plots, these trends were insignificant ($p>.10$). There was a significant negative relationship between butterfly abundance and cheatgrass ($p=.04$) as well as a highly significant relationship between butterfly species richness and cheatgrass cover of plots ($p=.001$). These results suggest that cheatgrass infestation in foothills grasslands may reduce populations of mammals, butterflies, and possibly birds. Cheatgrass should be considered when managing wildlife and wildlife habitats.

Keywords: Cheatgrass, Foothills Grassland, Mammals, Birds, Butterflies, Rocky Mountains, Invasive Plants, Ground Nesting Birds, Grass-Dependent Butterflies, Remote Triggering Cameras, Wildlife

Introduction

This study aims to give a preliminary indication as to whether cheatgrass affects distributions and populations of mammals, birds, and butterflies. We hypothesized that there would be lesser numbers and lesser species richness (total number of species) of native mammals, birds, and butterflies in areas heavily infested with cheatgrass. Testing this hypothesis is critical to good wildlife management and strategic planning of cheatgrass removal, especially in sensitive wildlife habitat. Few studies have investigated the relationships between cheatgrass and large mammals, birds, or butterflies.

Studies about the effect of cheatgrass on small mammals are abundant and the results are quite consistent: small mammals are negatively impacted by cheatgrass. Freedman et al. (2014) found that the abundance and diversity of small-mammal communities in the Great Basin decreased with increasing abundance of cheatgrass. Similarly, Ostoja and Schupp (2009) found that total rodent abundance was 6.1 times greater in sagebrush-dominated areas relative to cheatgrass-dominated areas in the Great Basin. Although countless studies replicated these findings for small mammals, to our knowledge no such studies have been conducted for larger mammals (ranging from lagomorphs to ungulates).

The impact of invasive plants on birds has been studied, but not specifically the impact of cheatgrass. Ortega et al. (2006) found that spotted knapweed reduced chipping sparrows' reproductive success and site fidelity, likely because of reduced food availability in areas infested by knapweed. Conversely, Scheiman et al. (2003) found that western meadowlark nesting success was positively associated with leafy spurge cover in North Dakota. These studies indicate that birds are sensitive to changes in the vegetative cover of their habitats, and therefore may be sensitive to cheatgrass.

101 Butterflies present another interesting gap in the scientific literature. Fleishman et al. (2005) concluded
102 that butterfly numbers within their study area in the Muddy River drainage (Nevada) were not impacted
103 by non-native plants but were influenced by nectar availability. Cheatgrass has the ability to form
104 monocultures that outcompete other plants (Young et al. 1987), including potential nectar sources,
105 indicating that cheatgrass may be detrimental to butterflies. Other research does cite invasive plants as
106 important stressors in butterfly communities (Keeley, 2017), but none of these studies looked specifically
107 at relationships between butterflies and cheatgrass.

108

109 To understand the relationships between cheatgrass and numbers of mammals, birds, and butterflies we
110 selected eight circular plots, each 75 meters in radius, containing varying amounts of cheatgrass cover.
111 We used a combination of remote-triggering cameras (for mammals), point counts (for birds), and transect
112 counts (for butterflies) to measure the number and species richness of animals on our plots.

113

114 **Study Area**

115

116 The study area lies at elevations between 1684-1790 m at the base of the Colorado Front Range foothills,
117 where the Great Plains abut the Southern Rocky Mountains. Elevation increases gradually from east to
118 west across the study area. Climate within this lower foothills region is characterized by relatively mild
119 but snowy winters and warm, dry summers. The mild winter temperatures result from frequent warm,
120 down-sloping winds, sometimes referred to as Chinooks. Annual precipitation averages 45-50 cm/year,
121 with approximately half of that amount falling during the March-June spring growing season (from US
122 Climate Data, 2020). As a result, maximum greening of native grasses and peak wildflower bloom usually
123 occur in May and June.

124

125 In early May, 2020, we established eight, 75 meter radius monitoring plots on the Boulder County Parks
126 and Open Space owned Pierce and Trevarton properties south of Lyons, Colorado in R 70W, T3N, S 31;
127 and R 70W, T 2N, S 6 (Figure 1, Table 1). Vegetation on these two properties consists primarily of
128 foothills mixed-grass prairie, foothills shrub, and ponderosa pine woodland (Baker and Galatowitsch
129 1985, Colorado Natural Areas Program 1998). Plant names that follow are from Wittmann and Weber
130 (2011).

131

132 In relatively flat areas with deep soils, the grassland tends to be dominated by green needlegrass (*Nassella*
133 *viridula*) and western wheatgrass (*Pascopyrum smithii*), with needle-and-thread grass (*Hesperostipa*
134 *comata*), junegrass (*Koeleria macrantha*), little bluestem (*Schizachyrium scoparium*), and non-natives
135 such as Canada bluegrass (*Poa canadensis*) and smooth brome (*Bromopsis inermis*) well represented.
136 Non-native cheatgrass (*Anisantha tectorum*) and crested wheatgrass (*Agropyron cristatum*) tend to thrive
137 in rocky or upland areas, along with native three-awn (*Aristida purpurea*) and buffalograss (*Buchloe*
138 *dactyloides*).

139

140 Flat-bottomed ravines cutting through the properties support extensive patches of native shrubs, including
141 skunkbrush (*Rhus trilobata*), wild plum (*Prunus americana*), mountain mahogany (*Cercocarpus*
142 *montanus*), chokecherry (*Padus virginiana*) snowberry (*Symphoricarpos* sp.), wild licorice (*Glycyrrhiza*
143 *lepidota*), and hawthorne (*Crataegus macracantha*). Scattered hackberry trees (*Celtis reticulata*) and box
144 elders (*Negundo aceroides*) grow in the relatively wet canyon bottoms, along with small patches of native
145 tallgrasses, including switchgrass (*Panicum virgatum*) and Indiangrass (*Sorghastrum avenaceum*).
146 Ponderosa pines (*Pinus ponderosa*) and Rocky Mountain junipers (*Sabina scopularum*) are scattered
147 throughout the study area, becoming most numerous in rocky uplands.

148

149 Plot 1 partially contains a one hectare prairie dog colony that extends beyond the plot to the east.
150 Vegetation over this colony consists primarily of cheatgrass and non-native forbs. Within the other seven
151 plots, we observed few obvious signs of disturbance by burrowing animal colonies or by human activities,
152 though a barely discernible two-track road cuts through portions of plots 7 and 8. A heavily-traveled two-
153 lane highway (U.S. 36) passes within 150 m of plots 1 and 5 (Figure 2). Barbed-wire fences border both
154 the Pierce and Trevarton properties, but none pass through our plots.

155

156 **Methods**

157

158 **Vegetation Sampling**

159

160 We located the eight circular, 75 m radius plots in areas of the Pierce and Trevarton properties where the
161 primary vegetation type is foothills grassland (Kuchler 1964). When we first established our plot locations
162 we tried to maximize the variability of cheatgrass cover in our plots by choosing 4 plots that appeared to
163 have high cheatgrass cover and 4 plots that appeared to have low cheatgrass cover.

164

165 To confirm our visual evaluation of the cheatgrass cover and to allow quantitative analysis we undertook
166 quantitative sampling of plot vegetation. To maximize the uniformity of our sampling (thereby yielding
167 better results), we used a Point Intercept Method with a modified distribution of sampling locations.

168 Standard radial vegetation sampling techniques (Figure 3a) are biased towards vegetation at the center of
169 the plot as they do not account for the non-linear increase of circle area with increased circle radius
170 (caused by the r^2 term in $A = \pi r^2$). We used a method that eliminated this bias, providing more
171 representative sampling, while only marginally increasing the time required to sample each plot.

172

Our method was as follows: we marked the center of each plot with a T-post. During the first two weeks of June we then laid out 8 radial transects 75 meters in length on each plot, originating at the center T-post and extending in each cardinal and inter-cardinal direction. To account for the bias described earlier, sampling locations were pushed outwards away from the center of the plot and closer to the outer edge of the plot (Figure 3b). The exact location of the samples was mathematically determined by splitting each plot into 5 sections of equal area delimited by 5 rings (with the 5th ring being the outer circumference of the plot). Each ring's radius was denoted as r_n with r_1 being the radius of the innermost ring. In order to make each section contain an equal area, the radius of each ring was calculated with the following equation: $r_n = \sqrt{n} * \left(r / \sqrt{n_{rings}} \right)$ where r is the radius of the plot (75 meters) and n_{rings} is the number of rings (5).

The 4 outer sampling locations on each transect were placed halfway between the nearest pair of rings. To further improve the method we rotated the second to innermost sampling location and the second to outermost sampling location by 22.5 degrees around the center post, thereby maximizing the distance between sampling locations (Figure 3c). Instead of laying out an additional 8 transects in order to locate these rotated sampling locations, we triangulated their positions on the 8 main radial transects such that a short sub-transect stretching between each pair of adjacent radial transects could be used to accurately locate the rotated sampling locations (Figure 4).

To locate the innermost sampling location on each transect, we divided the innermost (circular) ring of radius r_1 into two equal areas split by a ring of radius $r_1 / \sqrt{2}$. The innermost sampling locations for the four cardinal transects were placed at the midpoint between r_1 and $r_1 / \sqrt{2}$. The innermost sampling locations for the four inter-cardinal transects were placed at $r_1 / \sqrt{2}$, as moving them any further inward would excessively crowd them toward the center of the plot. As a result 4 of the 8 innermost samples were

197 located 29 meters from the center T-post and the other 4 were located 17 meters from the center T-post
198 (Figure 3d).

199

200 This method greatly increased the uniformity of our sampling on each plot, yielding results much more
201 representative of the actual vegetative cover of our plots while only marginally increasing the time
202 required to sample each plot. The distances for each sampling location from the center T-post can be
203 found in Table 2.

204

205 To sample the vegetation at each sampling location we used a Point-Intercept with Grid Quadrant Method
206 similar to the one described in Lutes et al. (2006). We used a 0.7 meter square sampling frame with 2 sets
207 of 5 strings stretched across each side at decimeter intervals, each set of strings positioned directly above
208 or below the other set. The frame was systematically placed at each sampling location. A metal pin was
209 lowered at each point where two strings of one set crossed (there were 25 such points), and the pin was
210 carefully aligned to the corresponding strings in the second set of strings below in order to ensure that the
211 pin was always at the right location and perpendicular to the frame's plane. We then noted whether the tip
212 of the pin was touching cheatgrass plants, cheatgrass litter, bare ground / rocks, non-cheatgrass litter, or
213 non-cheatgrass plants. The data for each pin were entered into a custom iPad app we had previously
214 developed. The app automatically synthesized and arranged the data by plot, making it exportable as a
215 CSV file. This app allowed us to streamline and accelerate the vegetation sampling process while
216 minimizing errors.

217

218 **Mammal Sampling**

219

220 We mounted a Stealth Cam G45NGX remote-triggering camera 0.9 meters off the ground on each T-post,
221 secured to a wooden block by zip-ties threaded through sets of holes in the block, such that the camera

could be oriented in each cardinal direction (Figure 5). The cameras were set to maximum sensitivity with a 15 second recovery time. We chose this short recovery time to ensure we did not miss mammals, while keeping false positive detections to a manageable level.

225

We began observation at 0000 MST on May 14th and ended observation at 2359 MST on August 13th. All cameras were originally pointed north. At intervals ranging from one to two weeks (on May 27th, June 14th, June 24th, July 11th, July 18th, July 24th, and August 7th) we rotated all the cameras clockwise by 90 degrees and replaced the batteries and the SD cards. In this manner all the cameras completed two observational rotations around their posts during the observation period. There was no malfunction or unexpected interruption in the cameras' operation over the course of the study.

232

We identified and counted all mammals in the images from the remote-triggering cameras without relying on previous or subsequent images for detection or identification of the mammals. Individual mammals were counted regardless of whether they had appeared in other images. Lingering mammals were effectively recounted roughly every 15 seconds when they remained in the trigger zone due to the 15 second recovery time, thus informally accounting for the duration of animal activity within the trigger zone. All the mammals we detected were categorized as one of the following: mule deer, white-tailed deer, unidentified deer, elk, unidentified ungulate, coyote, bobcat, prairie dog, unidentified cottontail, striped skunk, black bear, and unidentified mammal. When performing analysis we excluded mammals that were detected on camera rotation days, thereby eliminating data consistency problems resulting from mammals being detected while certain cameras were nonoperational.

243

244 **Breeding Bird Surveys**

245

246 We counted birds seen or heard from the center points of the eight monitoring plots during 8 minutes on
247 the mornings of 29 May, 16 and 29 June, and 9 July, beginning each survey at sunrise and completing
248 sampling of all eight plots by 0730 MST. We varied the order of the plot counts during each replication to
249 reduce temporal biases that might stem from sampling bird populations at varying times of the early
250 morning. As we entered each plot we counted any birds flushed from the vegetation. We noted numbers
251 of birds seen or heard perching or foraging within each monitoring plot but did not record numbers of
252 individuals flying over or through the plots without foraging (Ralph et al. 1998). Swallows were counted
253 when they were flying below the tops of the tallest trees and shrubs and their irregular movement patterns
254 suggested that they were foraging.

255

256 **Butterfly Surveys**

257

258 We counted butterflies seen along 150 m north / south transects bisecting each monitoring plot during a
259 time interval of 5 minutes on 6 and 20 June; 2, 16, and 27 July; and 7 August. We walked slowly (2 km /
260 hr) along each transect, using binoculars and cameras with telephoto lenses to identify all butterflies seen
261 within 30 m. Counts were carried out between 0745-0945 MST on calm (peak wind velocity ≤ 20 km/hr)
262 and clear (mean cloud cover $\leq 30\%$) mornings when the air temperature exceeded 18° C. Plot sampling
263 order was rotated during each replication to reduce temporal biases.

264

265 **Additional Data Collection**

266

267 In addition to measuring our primary variables we quantified other properties of our plots, notably slope,
268 distance to highway U.S. 36 (effectively the distance to the nearest human development), shrub / tree
269 cover, and percent bare and rocky ground. We used Google Earth Pro to approximate the first three

270 variables and data from our vegetation sampling to calculate the percentage of bare and rocky ground in
271 each plot.

272

273 To measure the slope of the plots we identified the highest and lowest points in each plot and divided the
274 difference in altitude of these two points by the distance between them. This gave us an approximation of
275 the slope of each the plot. To measure the distance to highway U.S. 36 we used the “measure” tool in
276 Google Earth Pro to measure the shortest distance between the center T-post of each plot (identified using
277 its GPS coordinates) and the nearest edge of highway U.S. 36. To calculate the shrub and tree cover we
278 used a 9 by 9 grid of points cropped into a circle containing 69 points. This was overlaid on the satellite
279 imagery such that it corresponded to the size of a plot, with points spaced by the equivalent of 15 m. For
280 each plot we identified whether each of the 69 points was over a tree, a shrub, or neither.

281

282 **Data Analysis**

283

284 The total cheatgrass cover of each of our plots was calculated by summing up the number of times our
285 vegetation sampling pin touched cheatgrass or cheatgrass litter and dividing the resulting value by the total
286 number of pin samples in each plot (1,000). We used this calculated estimate of cheatgrass cover
287 (essentially the density of cheatgrass plants and cheatgrass litter) in all our analyses.

288

289 We used the software package R to analyze our data with a combination of Linear Correlation analysis
290 (using Pearson Correlation Coefficients / Tests), as well as ANOVA analysis (using “aov” function in R).
291 We also verified our assumptions for these tests using the Shapiro-Wilk Normality Test and the Fligner-
292 Killeen Test of Homogeneity of Variances (when working with ANOVA analysis). For the data that failed
293 these tests we used the Kruskal-Wallis Rank Sum Test instead of the usual “aov” function in R.

294

295 For the ANOVA analysis we split our plots into two categories: low cheatgrass and high cheatgrass. We
296 defined any plot with less than 0.2 cheatgrass cover as low cheatgrass and any plot with more than 0.2
297 cheatgrass cover as high cheatgrass. This categorization gave us 4 high cheatgrass and 4 low cheatgrass
298 plots, as we had originally planned when selecting our plot locations. The cheatgrass cover of the low
299 cheatgrass plots varied from 0.031 cheatgrass cover to 0.135 cheatgrass cover, while the cheatgrass cover
300 of our high cheatgrass plots varied from 0.235 cheatgrass cover to 0.35 cheatgrass cover.

301

302 We used the R^2 value of both the Pearson and ANOVA tests to determine which best fit our data. We
303 generally found that our bird and mammal data best fit an ANOVA style model, while our butterfly data
304 clearly showed linear trends. We then consistently used the best fit model for our analysis, except for
305 visualization purposes. When our independent variable was not cheatgrass cover we always used Linear
306 Correlation Analysis.

307

308 **Results**

309

310 **Mammals**

311

312 During the 3 months of camera operation we recorded 18,326 images, 392 of which were identified as
313 containing mammals, for a total of 561 counted mammals (Table 3). We used only 530 mammals in our
314 analyses as the rest were detected on days when we rotated the cameras (see methods section). The most
315 frequently detected mammal species was mule deer (comprising 44% of detections), followed by elk (6%
316 of detections) and coyote (3% of detections). The cameras also detected significant quantities of
317 unidentified deer (29% of detections), unidentified ungulates (7% of detections), and unidentified
318 mammals (4% of detections). White-tailed deer, bobcat, cottontail species, and striped skunk represented
319 a combined total of 5% of detections. We detected only one black bear.

320

321 Scatterplots suggest a strong negative relationship between total mammals and cheatgrass cover (Figure
322 6) as well as a weaker negative relationship between mammal species richness (total number of species)
323 and cheatgrass cover (Figure 7). We then split all mammals into ungulates, carnivores / omnivores, and
324 lagomorphs / rodents, analyzing each category separately. Ungulates, like total mammals, showed a
325 relatively strong negative relationship with cheatgrass cover (Figure 8). Carnivores / omnivores were
326 weakly negatively correlated with cheatgrass cover (Figure 9), while lagomorphs / rodents showed a weak
327 positive relationship with cheatgrass cover (Figure 10).

328

329 Our ANOVA analysis showed greater numbers of mammals on low cheatgrass plots compared to high
330 cheatgrass plots ($F(1,6)=9.488$, $MSE=932$, $p=.02$, Figure 11a). We found no significant difference in
331 mammal species richness between low and high cheatgrass plots, but we did observe a weak negative
332 relationship ($F(1,6)=2.778$, $MSE=1.125$, $p=.15$, Figure 11b). A significant negative relationship was
333 found between ungulates and cheatgrass cover ($F(1,6)=8.783$, $MSE=834$, $p=.03$, Figure 11c), but only
334 insignificant relationships were found between carnivores / omnivores and cheatgrass cover ($\chi^2=0.35$,
335 $p=.55$, $df=1$, Figure 11d) and between lagomorphs / rodents and cheatgrass cover ($\chi^2=0.036$, $p=.85$,
336 $df=1$, Figure 11e).

337

338 **Birds**

339

340 During four, 8-minute point-counts, we observed a total of 26 potential breeding species on the 8 plots
341 (Table 4). Lark sparrows were most abundant (10.25 / count), followed by spotted towhees (5.50 / count)
342 and western meadowlarks (4.00 / count). Obligate ground-nesters comprised 39.3% of all birds observed.
343 Shrub-nesters, tree cavity-nesters, cliff-nesters (mostly swallows), lower tree-canopy nesters, and habitat
344 generalists comprised most of the remaining observed individuals.

345

346 Scatter plots suggest a negative relationship between cheatgrass cover and mean birds per plot (Figure
347 12); and a negative relationship between cheatgrass cover and total bird species per plot (Figure 13).

348

349 Since ground-nesting birds may depend on native grass cover for nesting and foraging (Kelsey 2010,
350 Sheridan et. al. 2020), we expected to find a strong negative relationship between mean ground-nesting
351 birds and mean cheatgrass cover. However, ground-nesting bird results were similar to results for all bird
352 species (Figures 12 and 14). Extent of native grass cover is likely just one of several factors contributing
353 to ground-nesting bird abundance. We observed lark sparrows and western meadowlarks foraging
354 primarily in areas dominated by native grass cover, but we also observed individuals perching frequently
355 on shrubs and small trees, including in areas with cheatgrass (see Discussion section, below).

356

357 Our ANOVA analysis showed an insignificant negative relationship between mean birds per plot and
358 cheatgrass cover ($F(1,6)=3.375$, $MSE=1.333$, $p=.12$, Figure 15a). The negative relationship between mean
359 ground-nesting birds and cheatgrass cover was also insignificant ($\chi^2=4.0514$, $p=.12$, $df=1$, Figure 15b).
360 We found that a linear model best fit the correlation between bird species richness (total number of bird
361 species) and cheatgrass cover, showing an insignificant negative trend ($r=-0.60$, $p=.12$, Figure 13).

362

363 **Butterflies**

364

365 During six, 5-minute transect surveys on each plot, we observed a total of 26 butterfly species on the 8
366 plots (Table 5). See Appendix I for scientific names. Variegated fritillaries (89.67/survey), habitat
367 generalists that occasionally invade the Rocky Mountain region in large numbers in response to
368 environmental stresses in the southern United States (Opler 1999), comprised more than 62% of all
369 butterflies observed. Other numerous species, with host plants in parentheses, included clouded sulphur

(8.17 / survey; legumes), common checkered-skipper (7.50 / survey; mallows), and orange sulphur (5.50 / survey; legumes). We observed six species that lay their eggs on grasses and that might be particularly sensitive to cheatgrass invasion: common ringlet (0.17 / survey), common wood nymph (3.00 / survey), dark wood nymph (0.33 / survey), green skipper (0.17 / survey), Arogos skipper (0.67 / survey), and taxiles skipper (0.17 / survey).

375

Scatterplots suggest a strong negative relationship between cheatgrass cover and mean butterflies per plot (Figure 16); and a strong negative relationship between cheatgrass cover and total species per plot (Figure 17). The relationship for grass-dependent species appears weaker, due in part to the small number of individuals detected (Figure 18).

380

Our Linear Correlation Analysis showed a significant negative relationship between mean butterflies and cheatgrass cover ($r=-0.73$, $p=.04$, Figure 16) as well as an insignificant negative relationship between mean grass dependent butterflies and cheatgrass cover ($r=-0.63$, $p=.097$, Figure 18). A highly significant negative relationship was found between butterfly species richness (total number of butterfly species) and cheatgrass cover ($r=-0.92$, $p=.001$, Figure 17).

386

387

Discussion

388

Summary of Results

390

Results of this single-season study suggest that cheatgrass infestation may negatively impact populations of mammals, birds, and butterflies in foothills grasslands (Table 6). Cheatgrass cover seemed only weakly related to mammal and bird species richness but appeared to have a strong negative impact on butterfly species richness.

395

396 We were intrigued that ANOVA analysis was the best-fitting model for mammals and birds, whereas
397 Linear Correlation Analysis worked best for butterflies. We hypothesize that mammals and birds, due to
398 their large home ranges, are relatively unaffected by cheatgrass up to a certain threshold level above
399 which mammals and birds avoid areas with cheatgrass. As they forage, areas with sufficient desirable
400 plants may attract them even if the desirable plants are interspersed with some cheatgrass. However, once
401 cheatgrass cover reaches a certain threshold, the foraging in that particular area may become unattractive
402 and the mammals and birds avoid it. Conversely, butterflies have small home ranges and are dependent on
403 native grasses and forbs within a small area for reproduction and nectar. Therefore, any reduction in
404 native plant density, even within a relatively small area, may directly drive down butterfly populations.
405 Further research with larger data sets would help to confirm or reject this hypothesis.

406

407 **Additional Factors**

408

409 Although our study does not prove causal relationships, we do believe that the differences in animal
410 populations among plots were driven by cheatgrass infestation. This tentative conclusion is supported by
411 the generally weak relationships between observed animal populations and plot characteristics other than
412 cheatgrass cover.

413

414 Mean slopes varied among plots, from approximately 0.10 in Plots 1 and 5 to 0.28 in Plot 4. Figure 19
415 shows the relationship between mean slope of each plot and numbers of detected mammals, birds, and
416 butterflies. This figure shows only very weak relationships between slope and the number of mammals
417 ($r=-0.59$, $p=.13$) and butterflies ($r=-0.56$, $p=.15$). There was no discernible relationship between birds and
418 slope ($r=-0.23$, $p=.59$).

419

Shrub and tree cover within the plots varied from nearly 45% in Plot 2, to 4% in plot 5. Shrubs provide resting, hiding, and foraging habitat for deer and elk, the most frequently detected mammals, along with nesting, perching, and foraging habitat for a preponderance of the observed bird species. Whereas butterflies typically derive nectar from blooming wildflowers growing in grassy areas, several locally common butterfly species lay their eggs on shrub leaves or perch on shrubs while resting or patrolling breeding territories (Chu and Jones 2020). Figure 20 shows the relationship of shrub and tree cover within individual plots to numbers of detected mammals, birds, and butterflies. This figure indicates a weak positive relationship between shrub and tree cover and numbers of butterflies per plot ($r=0.63$, $p=.09$) but no discernible relationships between shrub and tree cover and numbers of mammals ($r=-0.24$, $p=.56$) or birds ($r=0.21$, $p=.62$).

430

We noticed that cheatgrass appeared to be more abundant in bare or rocky areas. Figure 21 shows the relationship between percent of bare and rocky ground within individual plots to numbers of detected mammals, birds, and butterflies. Here the scatterplots indicate a significant negative relationship between percent of bare and rocky ground and butterflies ($r=-0.72$, $p=.04$) but no discernible relationship between percent of bare and rocky ground and numbers of mammals ($r=-0.27$, $p=.51$) or birds ($r=0.13$, $p=.77$).

436

Noting that two of the plots lie within close proximity to a busy highway, we also compared numbers of detected mammals, birds, and butterflies to plot distance from the highway (Figure 22). We found very weak negative relationships between numbers of mammals ($r=-0.59$, $p=.12$) and butterflies ($r=-0.57$, $p=.14$) and distance to highway U.S. 36. No discernible relationship was found between birds and distance to the highway ($r=-0.22$, $p=.59$). This may simply be a result of the plots near the highway being generally lushier than our more remote plots, thus possibly providing more food and shelter for animals. No discernible relationship was found between birds and distance to the highway ($r=-0.22$, $p=.59$).

444

445 We suspect that the fences that border both the Pierce and Trevarton properties may contribute to patterns
446 of mammal distribution and movement, though their effects are difficult to quantify. Barbed wire fences
447 have been shown to restrict movements of large mammals (Gates et al. 2012).

448

449 A variety of studies have indicated that total volume of foliage can be a strong predictor of abundance of
450 both birds and butterflies (Degraaf et al. 1998, Lee et al. 2015, Mills et al. 1991). Cheatgrass infestation
451 may reduce the total volume of foliage by crowding out native grasses and forbs (Billings 1994,
452 Parkinson et al. 2013). Since cheatgrass is a winter annual that greens up in early spring and becomes
453 senescent by early summer, areas of infestation may become ecological deserts that provide neither
454 breeding habitat nor forage for most birds and butterflies.

455

456 In contrast, native shrub cover increases the availability of breeding niches while also providing an
457 additional food source for grassland birds and butterflies. Higher density of native grasses, native forbs,
458 and native shrubs may partially explain why our lower-elevation plots supported higher numbers of
459 mammals, birds, and butterflies.

460

461 **Future Research**

462

463 Future research might include small mammal trapping and insect netting. As these animals typically have
464 smaller home ranges than large mammals and birds and also depend on native grasses and forbs for food
465 and shelter, we would expect their populations to be negatively impacted by cheatgrass cover. Clipping
466 and weighing of grasses and forbs within nested plots would help to determine the relationship of total
467 volume and weight of native and non-native vegetation to numbers of mammals, birds, and butterflies.
468 Longitudinal studies of plots with comparable amounts of tree and shrub cover would give us a clearer
469 view of the direct impact of cheatgrass cover on wildlife populations.

470

471 Future research should investigate the impact on wildlife of different treatment methods to control
472 cheatgrass. We recommend that such research compare wildlife populations over appropriately long
473 timescales on untreated plots relatively free of cheatgrass, on untreated plots infested by cheatgrass, and
474 on previously infested treated plots. The standard for treatment success should include restoration to
475 conditions similar to those found in untreated, relatively cheatgrass free areas, not just improvement over
476 untreated cheatgrass infested areas. A failure to investigate the effect of cheatgrass treatment methods on
477 wildlife could potentially cause the inadvertent degradation of valuable wildlife habitat. For example,
478 some herbicides are known to negatively impact certain butterfly species (Russell and Schultz 2010).
479 Without studies investigating the effects of herbicides on butterflies, herbicides might be used when
480 alternative methods, such as those suggested by Blumenthal et al. (2010), would be more appropriate.

481

482 In October 2020 the Calwood fire burned through four of our study plots. A follow-up study,
483 documenting observed fire impacts on cheatgrass cover, native plant cover, and wildlife populations,
484 would offer a rare opportunity to explore the role that fire plays in cheatgrass growth and in the
485 regeneration of native foothills grassland ecosystems.

486

487 **Acknowledgments**

488

489 Timothy Seastedt provided bountiful advice, knowledge, encouragement, and positivity that made this
490 research project possible. Alex Markevich shared wonderful insights and kindly proofread our paper.
491 Carron Meaney generously contributed her expertise on mammals and offered to coordinate field work.
492 Lynn Riedel patiently informed and educated us on vegetation sampling, providing the knowledge we
493 needed to design this project. Susan Spaulding and Steve Sauer helped us with this project from start to

494 finish and provided a vital link to Boulder County Parks and Open Space. And thank you to Boulder
495 County Parks and Open Space and Boulder County Nature Association for funding this research project.

496

497 **Literature Cited**

498

499 Baker, W. L., and Galatowitsch, S. M. (1985), The Boulder tallgrass prairies. *Boulder County Nature*
500 *Association Publication*, (3).

501

502 Billings, W. D. (1994), Ecological impacts of cheatgrass and resultant fire on ecosystems in the western
503 Great Basin. *Proceedings–Ecology and Management of Annual Rangelands*’. (Eds SB Monsen, SG
504 Kitchen) pp, 22-30.

505

506 Blumenthal, D. M., Norton, A. P., and Seastedt, T. R. (2010), Restoring competitors and natural enemies
507 for long-term control of plant invaders. *Rangelands*, 32(1), 16-20.

508

509 Chu, J. R., and Jones, S. R. (2020). Butterflies of the Colorado Front Range. Boulder County Nature
510 Association, Boulder, CO.

511

512 Colorado Natural Areas Program. (1998), *Native Plant Revegetation Guide for Colorado*. BiblioGov
513 Project. <https://cpw.state.co.us/Documents/CNAP/RevegetationGuide.pdf>.

514

515 DeGraaf, R. M., Hestbeck, J. B., and Yamasaki, M. (1998), Associations between breeding bird
516 abundance and stand structure in the White Mountains, New Hampshire and Maine, USA. *Forest Ecology*
517 *and Management*, 103: 217-233., 103.

518

- 519 Fleishman, E., Mac Nally, R., and Murphy, D. D. (2005), Relationships among non-native plants,
520 diversity of plants and butterflies, and adequacy of spatial sampling. *Biological Journal of the Linnean*
521 *Society*, 85(2), 157-166.
- 522
- 523 Freeman, E. D., Sharp, T. R., Larsen, R. T., Knight, R. N., Slater, S. J., and McMillan, B. R. (2014),
524 Negative effects of an exotic grass invasion on small-mammal communities. *PLoS One*, 9(9), e108843.
- 525
- 526 Gates, C. C., Jones, P., Sutor, M., Jakes, A., Boyce, M. S., Kunkel, K., and Wilson, K. (2012), The
527 influence of land use and fences on habitat effectiveness, movements and distribution of pronghorn in the
528 grasslands of North America. In *Fencing for conservation* (pp. 277-294). Springer, New York, NY.
- 529
- 530 Keeley, W. (2017), Butterfly community monitoring on City of Boulder Open Space and Mountain Parks
531 property: analysis of the changes in species richness and diversity from 2002 to 2016. *Publications-*
532 *Colorado Natural Heritage Program*.
- 533
- 534 Kelsey, R. (2010) Enhancing grassland restoration for grassland birds. *Grasslands* (Winter 2010, 9-14).
- 535
- 536 Kuchler, A. M. (1964), Potential natural vegetation of the conterminous United States. *Am. American*
537 *Geographical Society Publication*, 36.
- 538
- 539 Lee, C. M., Park, J. W., Kwon, T. S., Kim, S. S., Ryu, J. W., Jung, S. J., and Lee, S. K. (2015), Diversity
540 and density of butterfly communities in urban green areas: an analytical approach using GIS. *Zoological*
541 *Studies*, 54(1), 4.
- 542

- 543 Lutes, D. C., Keane, R. E., Caratti, J. F., Key, C. H., Benson, N. C., Sutherland, S., and Gangi, L. J.
544 (2006), FIREMON: Fire effects monitoring and inventory system. Gen. Tech. Rep. RMRS-GTR-164.
545 Fort Collins, CO: US Department of Agriculture, Forest Service, Rocky Mountain Research Station. 1
546 CD., 164.
547
- 548 Mills, G. S., Dunning Jr, J. B., and Bates, J. M. (1991), The relationship between breeding bird density
549 and vegetation volume. *The Wilson Bulletin*, 468-479.
550
- 551 Opler, P. A. (1999), A field guide to western butterflies. Houghton Mifflin Harcourt.
552
- 553 Ortega, Y. K., McKelvey, K. S., and Six, D. L. (2006), Invasion of an exotic forb impacts reproductive
554 success and site fidelity of a migratory songbird. *Oecologia*, 149(2), 340-351.
555
- 556 Ostoja, S.M. and Schupp, E.W. (2009), Conversion of sagebrush shrublands to exotic annual grasslands
557 negatively impacts small mammal communities. *Diversity and Distributions*, 15: 863-870. doi:10.1111/
558 j.1472-4642.2009.00593.x
559
- 560 Parkinson, H., Zabinski, C., and Shaw, N. (2013), Impact of native grasses and cheatgrass (*Bromus*
561 *tectorum*) on Great Basin forb seedling growth. *Rangeland Ecology & Management*, 66(2), 174-180.
562
- 563 Ralph, C. J., Sauer, J. R., and Droege, S. (Eds.). (1998), Monitoring bird populations by point counts.
564 DIANE Publishing.
565
- 566 Russell, C., and Schultz, C. B. (2010), Effects of grass-specific herbicides on butterflies: an experimental
567 investigation to advance conservation efforts. *Journal of Insect Conservation*, 14(1), 53-63.

568

569 Scheiman, D. M., Bollinger, E. K., and Johnson, D. H. (2003), Effects of leafy spurge infestation on
570 grassland birds. *The Journal of wildlife management*, 115-121.

571

572 Sheridan, K., Monaghan, J., Tierney, T. D., Doyle, S., Tweney, C., Redpath, S. M., and McMahon, B. J.
573 (2020), The influence of habitat edge on a ground nesting bird species: hen harrier *Circus*
574 *cyaneus*. *Wildlife Biology*, 2020(2).

575

576 Wittmann, R. C., and Weber, W. A. (2011), *Colorado Flora: Eastern Slope*. University Press of Colorado.

577

578 Young, J. A., Evans, R. A., Eckert, R. E., and Kay, B. L. (1987), Cheatgrass. *Rangelands Archives*, 9(6),
579 266-270.

580

581

582

583

584

585

586

587

588

589

590

591

592

593

Tables

594

595 Table 1. Plot information

596

	1	2	3	4	5	6	7	8
GPS Coordinates NS	40°9'56.90 ''N	40°10'4.2 0''N	40°10'9.1 0''N	40°10'18. 60''N	40°10'35.2 0''N	40°10'33.7 0''N	40°10'47.0 0''N	40°10'54.5 0''N
GPS Coordinates EW	105°16'12. 50''W	105°16'8. 50''W	105°16'7. 90''W	105°16'6. 50''W	105°15'35. 50''W	105°15'41. 80''W	105°15'34. 30''W	105°15'33. 30''W
Property	Pierce	Pierce	Pierce	Trevarton	Trevarton	Trevarton	Trevarton	Trevarton
Altitude (m)	1743	1726	1727	1747	1690	1699	1742	1773
Proximity to U.S. 36 (m)	114	185	249	373	130	188	389	484
Slope	0.102	0.108	0.153	0.275	0.102	0.140	0.182	0.229
Cheatgrass Cover	0.35	0.031	0.061	0.312	0.087	0.135	0.349	0.235
Shrub And Tree Cover	0.116	0.449	0.087	0.188	0.043	0.116	0.174	0.188

597

598

599

600

601

602

603

604

605

606 Table 2. Distances of sampling locations from center post

607

Sample Number	Samples on N, E, S, and W Transects	Samples on NE, SE, SW, and NW Transects	Rotated Sample
1	17 meters	29 meters	No
2	40 meters	40 meters	Yes
3	53 meters	53 meters	No
4	63 meters	63 meters	Yes
5	71 meters	71 meters	No

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626 Table 3. Total numbers of mammals detected by remote-triggering cameras mounted in the center of
 627 monitoring plots

628

Plot	1	2	3	4	5	6	7	8	Total
Mule Deer	8	27	23	19	65	57	34	2	235
White Tailed Deer	0	2	1	0	0	2	0	0	5
Misc. Deer	6	24	24	6	65	11	16	2	154
Elk	2	7	14	1	4	0	2	3	33
Ungulate	1	9	5	4	9	5	1	5	39
Coyote	6	1	3	0	8	3	0	0	21
Bobcat	0	1	0	1	1	1	0	2	6
Prairie Dog	3	0	0	0	0	0	0	0	3
Cottontail Species	3	0	0	0	0	3	0	0	6
Striped Skunk	0	0	0	2	5	0	0	0	7
Black Bear	0	0	0	0	0	0	0	1	1
Misc. Mammal	0	8	5	0	3	2	0	2	20
Total	29	79	75	33	160	84	53	17	530

629

630

631

632

633

634

635

636

637

638

639

640

641

642 Table 4. Mean number of birds seen or heard within monitoring plots during four, 8-minute point-counts

643

Species	1	2	3	4	5	6	7	8	Total
Broad-tailed Hummingbird		0.75	0.25	0.75			0.5	0.25	2.50
Mourning Dove		0.5			0.25				0.75
Northern Flicker					0.75				0.75
Western Wood-Pewee								0.25	0.25
Plumbeous Vireo								0.25	0.25
Blue Jay		0.25		0.25					0.50
Black-billed Magpie				0.25		1.50			1.50
Barn Swallow						0.25			0.25
Cliff Swallow			0.25		0.25	0.5	0.25		1.25
Black-capped Chickadee		0.25							0.25
White-breasted Nuthatch								0.25	0.25
Rock Wren				0.25					0.25
American Robin	0.25	0.25		0.25	0.5				1.25
Gray Catbird					0.25				0.25
Brown Thrasher	0.25								0.25
European Starling					0.25				0.25
Lesser Goldfinch			0.25						0.25
American Goldfinch	0.25				0.25	0.25			0.75
Lark Sparrow	1.0	0.75	1.75	0.5	1.75	2.00	1.25	1.25	7.75
Spotted Towhee	0.5	0.5		0.5	0.25	0.5	1.75	1.5	5.50
Yellow-breasted Chat		1.25	0.25		0.25				1.75
Western Meadowlark	0.25	0.5	0.5	0.5	1.25	1.0			4.00
Bullock's Oriole	0.25	0.25							0.50
Brown-headed Cowbird		0.25							0.25
Blue Grosbeak						0.25			0.25
Lazuli Bunting		0.25		1.0		0.25		0.5	2.00
Passerine species							0.5		0.50
Total	2.75	5.75	3.25	4.25	6.00	6.50	4.25	4.25	

644 Table 5. Mean number of butterflies observed within monitoring plots during six, 5-minute transect

645 surveys

646

Species	1	2	3	4	5	6	7	8	Total
Two-tailed Swallowtail	0.17						0.33	0.17	0.67
W. Tiger Swallowtail								0.17	0.17
Checkered White	0.17	1.17	0.67	0.17	0.17	0.17	0.67	0.33	3.50
Western White	1.50	0.67	0.17	0.17	1.17	0.67	0.17	0.33	5.50
Cabbage White		1.17	0.17		0.33				1.67
White Species	0.17	0.17	0.33	0.50	1.17				2.33
Clouded Sulphur	1.83	2.67	2.33		0.17	1.00		0.17	8.17
Orange Sulphur	0.33	2.33	1.83		0.83	0.17			5.50
Dainty Sulphur		0.17			0.17	0.17			0.50
Sulphur species		0.67							0.67
Elfin species		0.33							0.33
Melissa Blue			1.33						1.33
Reakirt's Blue								0.17	0.17
Gray Copper					0.17				0.17
Gorgone Checkerspot		0.17							0.17
Red Admiral		0.17							0.17
Variegated Fritillary	11.50	25.83	9.83	9.17	9.83	14.17	4.67	4.67	88.67
Speyeria Fritillary ¹	0.50	3.17	1.00	1.50	0.83	1.00	0.33		8.33
Hackberry Emperor		0.33	0.17	0.17	0.50	0.33	0.17		1.67
Milbert's Tortoiseshell			0.33						0.33
Common Ringlet				0.17					0.17
Common Wood Nymph		0.83	1.00	0.67	0.33	0.17			3.00
Dark Wood Nymph				0.17		0.17			0.33
Silver-spotted Skipper					0.50				0.50
Common Checkered-Skipper	0.17	1.33	3.67		1.00	0.67		0.67	7.50
Green Skipper		0.17							0.17
Arogos Skipper		0.33			0.17	0.17			0.67
Taxiles Skipper							0.17		0.17
Unidentified	0.17			0.17					0.33
Total	16.50	41.67	17.83	12.83	17.33	18.83	6.50	7.33	

647


1 Includes both “northwestern” and afrodite fritillary

Table 6. Summary of Results


Plot	1	2	3	4	5	6	7	8
Mammals	29	79	75	33	160	84	53	17
Mammal Species	5	5	4	4	5	6	2	4
Mean Birds	2.75	5.75	3.25	4.25	6.00	6.50	4.25	4.25
Bird Species	7	12	6	9	11	9	4	7
Mean Butterflies	16.50	41.67	17.83	12.83	17.33	18.83	6.50	7.33
Butterfly Species	8	17	12	9	15	12	7	9

Figures


Figure 1. Location of 8 plots within Open Space properties, with green pins indicating low cheatgrass plots and red pins indicating high cheatgrass plots (see Methods section)




668 Figure 2. Satellite map of 8 plots with green pins indicating low cheatgrass plots and red pins
669 indicating high cheatgrass plots (see Methods section)



672 Figure 3. Different ways to arrange sampling locations along radial transects within circular plots



675 Figure 4. Positioning of rotated sampling locations (black dots are sampling locations, grey dots
676 are flagging locations, black lines are radial transects, and grey lines are sub-transects)



678

679

680

681

682


683

684

685 Figure 5. Stealth Cam G45NGX remote-triggering camera mounted on T-post using a wooden
686 block and zip-ties



Figure 6. Total mammals and cheatgrass cover (showing line of best fit and 95% confidence interval)



721 Figure 7. Mammal species richness (total number of species) and cheatgrass cover (showing line
722 of best fit and 95% confidence interval)




Figure 8. Total ungulates and cheatgrass cover (showing line of best fit and 95% confidence interval)




Figure 9. Total carnivores / omnivores and cheatgrass cover (showing line of best fit and 95% confidence interval)






Figure 10. Total lagomorphs / rodents and cheatgrass (showing line of best fit and 95% confidence interval)



793 Figure 11. Bar graphs of mammals and cheatgrass (showing mean and standard deviation)

794



795 Figure 12. Mean birds and cheatgrass cover (showing line of best fit and 95% confidence
796 interval)





Figure 13. Bird species richness (total number of species) and cheatgrass cover (showing line of best fit and 95% confidence interval)



831 Figure 14. Mean ground nesting birds and cheatgrass cover (showing line of best fit and 95%
832 confidence interval)





Figure 15. Bar graphs of birds and cheatgrass (showing mean and standard deviation)

a. Mean Birds and Cheatgrass Cover



b. Mean Ground-Nesting Birds and Cheatgrass Cover




Figure 16. Mean butterflies and cheatgrass cover (showing line of best fit and 95% confidence interval)




Figure 17. Butterfly species richness (total number of species) and cheatgrass cover (showing line of best fit and 95% confidence interval)




Figure 18. Mean grass dependent butterflies and cheatgrass cover (showing line of best fit and 95% confidence interval)




Figure 19. Graphs of animals and plot slope (showing line of best fit and 95% confidence interval)




Figure 20. Graphs of animals and shrub / tree cover of plots (showing line of best fit and 95% confidence interval)




Figure 21. Graphs of animals and bare ground / rock cover of plots (showing line of best fit and 95% confidence interval)





Figure 22. Graphs of animals and plot distance to highway U.S. 36 (showing line of best fit and 95% confidence interval)



Appendix I. Scientific Names of Mammals, Birds, and Butterflies Mentioned in Text

Mammals

<u>Common Name</u>	<u>Scientific Name</u>
Mule Deer	<i>Odocoileus hemionus</i>
White-Tailed Deer	<i>Odocoileus virginianus</i>
Deer Species	<i>Odocoileus spp.</i>
Elk	<i>Cervus canadensis</i>
Ungulate	<i>Ungulata spp.</i>
Coyote	<i>Canis latrans</i>
Bobcat	<i>Lynx rufus</i>
Prairie Dog	<i>Cynomys ludovicianus</i>
Cottontail	<i>Leporidae spp.</i>
Striped Skunk	<i>Mephitis mephitis</i>
Black Bear	<i>Ursus americanus</i>
Unidentified Mammal	<i>Mammalia</i>

Birds

<u>Common Name</u>	<u>Scientific Name</u>
Mourning Dove	<i>Zenaida macroura</i>
Common Nighthawk	<i>Chordeiles minor</i>
Broad-tailed Hummingbird	<i>Selasphorus platycercus</i>
Turkey Vulture	<i>Cathartes aura</i>
Sharp-shinned Hawk	<i>Accipiter striatus</i>
Red-tailed Hawk	<i>Buteo jamaicensis</i>
Great Horned Owl	<i>Bubo virginianus</i>
Hairy Woodpecker	<i>Dryobates villosus</i>
Northern Flicker	<i>Colaptes auratus</i>
American Kestrel	<i>Falco sparverius</i>
Western Kingbird	<i>Tyrannus verticalis</i>
Say's Phoebe	<i>Sayornis saya</i>
Western Wood-Pewee	<i>Contopus sordidulus</i>
Plumbeous Vireo	<i>Vireo plumbeus</i>
Steller's Jay	<i>Cyanocitta stelleri</i>
Blue Jay	<i>Cyanocitta cristata</i>
Black-billed Magpie	<i>Pica hudsonia</i>
American Crow	<i>Corvus brachyrhynchos</i>
Tree Swallow	<i>Tachycineta bicolor</i>
Barn Swallow	<i>Hirundo rustica</i>
Cliff Swallow	<i>Petrochelidon pyrrhonota</i>

<u>Common Name</u>	<u>Scientific Name</u>
Black-capped Chickadee	<i>Poecile atricapilla</i>
White-breasted Nuthatch	<i>Sitta carolinensis</i>
Rock Wren	<i>Salpinctes obsoletus</i>
Townsend's Solitaire	<i>Myadestes townsendi</i>
American Robin	<i>Turdus migratorius</i>
Gray Catbird	<i>Dumetella carolinensis</i>
Brown Thrasher	<i>Toxostoma rufum</i>
Lesser Goldfinch	<i>Spinus psaltria</i>
American Goldfinch	<i>Spinus tristis</i>
Lark Sparrow	<i>Chondestes grammacus</i>
Chipping Sparrow	<i>Spizella passerina</i>
Green-tailed Towhee	<i>Pipilo chlorurus</i>
Spotted Towhee	<i>Pipilo maculatus</i>
Yellow-breasted Chat	<i>Icteria virens</i>
Western Meadowlark	<i>Sturnella neglecta</i>
Bullock's Oriole	<i>Icterus bullockii</i>
Brown-headed Cowbird	<i>Molothrus ater</i>
Common Grackle	<i>Quiscalus quiscula</i>
Virginia's Warbler	<i>Leiothlypis virginiae</i>
Yellow Warbler	<i>Setophaga petechia</i>
Black-headed Grosbeak	<i>Pheucticus melanocephalus</i>
Blue Grosbeak	<i>Passerina caerulea</i>
Lazuli Bunting	<i>Passerina amoena</i>

70
71

72 **Butterflies**

73

<u>Common Name</u>	<u>Scientific Name</u>
Western Tiger Swallowtail	<i>Papilio rutulus</i>
Two-tailed Swallowtail	<i>Papilio multicaudata</i>
Checkered White	<i>Pontia protodice</i>
Western White	<i>Pontia occidentalis</i>
Cabbage White	<i>Pieris rapae</i>
Clouded Sulphur	<i>Colias philodice</i>
Orange Sulphur	<i>Colias eurytheme</i>
Dainty Sulphur	<i>Nathalis iole</i>
Elfin species	<i>Callophrys sp.</i>
Gray Copper	<i>Lycaena dione</i>
Reakirt's Blue	<i>Echinargus isola</i>
Melissa Blue	<i>Plebejus melissa</i>
Rocky Mtn. Dotted Blue	<i>Euphilotes ancilla</i>
Monarch	<i>Danaus plexippus</i>
Variegated Fritillary	<i>Euptoieta claudia</i>
Northwestern Fritillary	<i>Speyeria aphrodite</i>

92	<u>Common Name</u>	<u>Scientific Name</u>
93	Aphrodite Fritillary	<i>Speyeria hesperis</i>
94	Coronis Fritillary	<i>Speyeria coronis</i>
95	Northern Crescent	<i>Phyciodes cocyta</i>
96	Pearl Crescent	<i>Phyciodes pallida</i>
97	Northern Checkerspot	<i>Chlosyne palla</i>
98	Gorgone Checkerspot	<i>Chlosyne gorgone</i>
99	Milbert's Tortoiseshell	<i>Aglais milberti</i>
100	Red Admiral	<i>Vanessa atalanta</i>
101	Painted Lady	<i>Vanessa cardui</i>
102	Hackberry Emperor	<i>Asterocampa celtis</i>
103	Common (Ocher) Ringlet	<i>Coenonympha tullia</i>
104	Common Wood Nymph	<i>Cercyonis pegala</i>
105	Dark Wood Nymph	<i>Cercyonis oetus</i>
106	Silver-Spotted Skipper	<i>Epargyreus clarus</i>
107	Common Checkered-Skipper	<i>Burnsius communis</i>
108	Green Skipper	<i>Hesperia viridis</i>
109	Arogos Skipper	<i>Atrytone arogos</i>
110	Taxiles Skipper	<i>Poanes taxiles</i>

111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133

Appendix II: Photos of Study Area, Monitoring Plots, and Selected Wildlife



Foothills grassland, with green needlegrass in foreground, Pierce Property. 5 June.



139
140 Western wheatgrass on Plot 5, Trevarton property, 16 June.
141



142
143 Prairie dog colony with cheatgrass and non-native forb cover on Pierce Property
144



Vegetation sampling on Plot 3, 4 June.



Plot 1, 24 June.



151
152
153

Plot 2, 24 June. Note shallow ravines with mountain mahogany and skunkbrush.



154
155
156

Plot 3, 24 June.



157
158
159

Plot 4, 24 June. Note extensive cheatgrass cover throughout.



160
161
162

Plot 5, 24 June



163
164
165

Plot 6, 24 June. Note shallow ravine with box elder and hackberry trees.



166
167
168

Plot 7, 24 June. Note cheatgrass cover, foreground, and ponderosa pine woodland, background.



169
170 Plot 8, 24 June. Note ponderosa pine woodland in background.
171



172
173 Hackberry emperor on snowberry bush in ravine near Plot 5.
174




175
176 Wood nymph resting on skunkbrush, Plot 2.
177



178
179 Bobcat, Plot 8.
180



Elk, Plot 2.



Young mule deer buck prancing through western wheatgrass, Plot 5.



White-tailed or hybrid deer in skunkbrush and mountain mahogany thicket, Plot 2, 4 June.

187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212

Appendix III. Github Repository of Cheatgrass Sampling App

213
214

215 The code for the iPad app we used to sample cheatgrass can be found here:

216 <https://github.com/gallus-gallus/Plot-Helper>

217

218 Current functionality may be limited. Questions should be addressed to:

219 anymmarkevich@gmail.com